



NUTRITIONAL STATUS OF PREGNANT WOMEN AND HISTORY PARITY FOR STUNTING IN TODDLER AGES 24-59

Lidyani Margareth, Nurul Husnul Lail*, Lisa Trina Arlym

Universitas Nasional, Jl. Sawo Mania No.61, Pejaten Barat, Ps. Minggu, Jakarta Selatan, Jakarta 12520, Indonesia

*nurulhusnul@civitas.unas.ac.id

ABSTRACT

Stunting is a condition where a child experiences growth disorders, which results in the child's height not being appropriate for his age as a result of chronic nutritional problems, namely a lack of adequate nutritional intake over a long period of time. Objective to analyze the relationship between maternal risk factors and the incidence of stunting among toddlers in the Karawaci Health Center working area. This research design uses an observational analytical quantitative research type with a case control design. The case samples in this study were toddlers aged 24-59 months who experienced stunting with a TB/U toddler Z score < -2 SD and the data was recorded at the Karawaci Health Center in 2023 as many as 123 toddlers, and a control sample of 123 toddlers. Data were analyzed using the chi square statistical test. Results of bivariate analysis Overall, of the 11 variables carried out in the research, 5 of them showed a relationship between these variables and the incidence of stunting in toddlers, including a relationship between parity, of stunting, in toddlers aged 24-59 months who are in the working area of the Karawaci Health Center. There is no relationship between the nutritional status of pregnant women, with the incidence of stunting in toddlers aged 24-59 months in the working area of the Karawaci Health Center. And the factor that poses the greatest risk for stunting is the age of the mother at risk during pregnancy, where stunting is 3.42 times more likely to occur than mothers of normal age during pregnancy. There is a relationship between the number of parities, maternal age at pregnancy, history of anemia during pregnancy, birth interval and maternal education level with the incidence of stunting.

Keywords: history of parity; nutritional status of pregnant women; stunting

How to cite (in APA style)

Margareth, L., Lail, N. H., & Arlym, L. T. (2025). Nutritional Status of Pregnant Women and History Parity for Stunting in Toddler Ages 24-59. *Indonesian Journal of Global Health Research*, 7(4), 1171-1176. <https://doi.org/10.37287/ijghr.v7i4.6843>.

INTRODUCTION

World Health Organization (WHO) and the World Bank Group, Stunting is estimated to affect 22% or 149.2 million children under 5 years globally in 2020. This figure has increased from 2019 which shows that stunting globally affects 21.3% or 144 million children. This happened because the impact of the pandemic occurred gradually, thus having an impact on food security and the community's economy. Therefore, an increase in the prevalence of stunting may be seen among children born in the first year of the pandemic. In 2020, more than half of the global percentage of children under 5 years experiencing stunting were in the Asian region with the highest rate of stunting cases at 53%, followed by the African region with 41%. The largest cases were found in the Asian region, namely 79 million cases. Southeast Asia has the second highest ranking for stunting cases, namely 15.3 million, below South Asia which reaches 54.3 million cases. Indonesia is the second country with the highest stunting cases in the Southeast Asia region after Timor Leste. Indonesia is still included in the category of very high stunting cases with a percentage of 31.8% in 2020 (Ministry of Health of the Republic of Indonesia, 2021).

Banten Province is among the top five regions with the highest stunting rates in Indonesia and is also included in 12 priority regions, according to the Indonesian Toddler Nutrition Status Survey (SSGBI) in 2021. Where there is not a single district/city where the prevalence is

below 10%. Based on data from the Indonesian Nutrition Status Survey (SSGI), the stunting rate in Banten in 2022 fell by 4.5% (24.5% in 2021 to 20% in 2022 below the national figure of 21.6%). However, there are still 5 districts or cities with a prevalence above the average rate of stunting under five in Banten. The yellow zone is in Cilegon City and Serang City, Lebak Regency, Tangerang Regency; while green is in South Tangerang and Tangerang City. Therefore, the Banten Provincial Government has committed to accelerating the prevention of stunting in Banten Province through cross-sectoral coordination, and strengthening through budgeting which must have clear output and contribute to reducing stunting. (Annur CM, 2023). Stunting incidents occur during pregnancy as a result of lack of nutrition during pregnancy, early initiation of breastfeeding in less than 1 hour of birth or not at all, breastfeeding stopping for approximately 6 months and the frequency of breastfeeding not being long enough, as well as providing food that accompanies breast milk for approximately 6-12 months, and the food given does not vary with frequency and texture that is not appropriate to age (Anggryni et al, 2021).

METHOD

This research design uses a type of quantitative analytical observational research with a case control design, which is an observational study that assesses the relationship between exposure and disease by determining a group of people who are sick (cases) and a group of people who are not sick (control) and then comparing the frequency of exposure in the two groups. The data collection technique uses purposive sampling. The population in this study were mothers who had children aged 24-59 months, as many as 991. The sample in this study was 123 cases of stunted toddlers and 123 who were not stunted as controls. Analisy with Chi-Square.

RESULT

Table 1.

Maternal characteristics based on nutritional status of pregnant women, history of parity in the of *stunting* among toddlers

Karakteristik	Frekuensi	Persentase (%)
<i>Stunting</i>	123	50
<i>Not stunting</i>	123	50
Nutritional Statys on Pregnancy		
KEK (LiLA < 23.5 cm)	70	28.4
Normal (LiLA ≥ 23.5 cm)	176	71.6
Parity		
Multipara	136	55.3
Primipara	110	44.7

Based on the distribution of maternal nutritional status during pregnancy, the results showed that of pregnant women with KEK nutritional status (LiLA < 23.5 cm), 34 children were found to have stunted toddlers (27.6%) and 26 children who were not stunted (21.1%). Meanwhile, from mothers who did not have CED (LiLA > 23.5 cm), 89 children (72.4%) were stunted and 97 children (78.9%) were not stunted. Based on the distribution of maternal factors, namely history of parity, among multiparous mothers (second or more pregnancies), 81 children were found with stunting (65.9%) and 55 children (44.8%) without stunting. Meanwhile, primiparous mothers (first pregnancy) had 42 children with stunting (34.1%) and 68 children without stunting (55.2%).

Table 2.
 Hasil Analisis Bivariat Variabel Independen terhadap Kejadian *Stunting*

Variabel Independen	Kejadian <i>Stunting</i>						p-value	OR
	<i>Stunting</i>		Tidak <i>Stunting</i>		TOTAL			
	f	%	f	%	f	%		
Status Gizi Ibu Saat Hamil								
KEK (LiLA < 23.5 cm)	34	56.7	26	43.3	60	100	0.299	1.42 (0.793 - 2.561)
Normal (LiLA ≥ 23.5 cm)	89	47.8	97	52.2	186	100		
Paritas Ibu								
Multipara	81	59.6	55	40.4	136	100	0.001	2.38 (1.425 - 3.991)
Primipara	42	38.2	68	61.8	110	100		

Based on the table of bivariate analysis results between risk factor variables for the incidence of stunting, it was found that the nutritional status of mothers during pregnancy did not have a significant relationship with the incidence of stunting in toddlers ($p > 0.05$). Meanwhile, the maternal parity variable shows that birth status/mother's parity is related to the incidence of stunting in toddlers ($p < 0.05$). Mothers with multiparous status have a 2.38 times greater risk of having stunted toddlers compared to mothers with primiparous status ($p = 0.001$; OR = 2.38; CI = 0.79-2.56).

DISCUSSION

The relationship between maternal nutritional status during pregnancy and the incidence of stunting

Based on table 2, it can be seen that from pregnant women with CED, 34 (27.6%) children were stunted, while from pregnant women who did not have CED, 89 (72.4%) children were stunted. In other words, in this study, the results showed that stunted children were more common in pregnant women who did not have CED. And from the table of bivariate analysis results, it was found that maternal nutritional status during pregnancy did not have a significant relationship with the incidence of stunting in toddlers ($p > 0.05$). According to theory, the nutritional status of the mother before and during pregnancy can influence the nutritional status of the mother and baby. The mother's nutritional intake affects the growth and development of the fetus, because the fetus's nutritional needs originate from the mother. If the mother experiences malnutrition, it can cause various risks such as bleeding, abortion, stillbirth, low birth weight (LBW) babies, congenital abnormalities, mental retardation, and so on (Sulistyoningsih, 2011). When women experience malnutrition in the final trimester, they are at risk of giving birth to babies with low birth weight (< 2,500 grams) (Arisman, 2004).

This is in line with research conducted by Anna (2021) in the Jatinom Community Health Center working area, Klaten, where the results of calculations using Chi-Square correlation obtained a p value of 0.605, which means there is no relationship between the history of KEK pregnant women and the incidence of stunting at the age of 3-5 years. Likewise, research by Kristiana et al (2016) showed that a history of CED during pregnancy was not statistically related to the incidence of stunting in toddlers (OR=0.61, 95% CI=0.32-1.14). The research is different from that conducted by Ratna, et al (2020), where it was found that mothers who had LiLA < 23.5 cm were at risk of giving birth to toddlers who experienced stunting ($p = 0.000$).

The relationship between maternal parity and the incidence of stunting

Based on the research results in table 2, it was found that from multiparous mothers there were 81 (65.9%) stunted toddlers, while from primiparous mothers there were 42 (43.1%) stunted toddlers. And the conclusion from the bivariate analysis was that birth status/mother's parity was related to the incidence of stunting in toddlers ($p < 0.05$). Mothers with multiparous status have a 2.38 times greater risk of having stunted toddlers compared to mothers with primiparous status ($p = 0.001$; OR = 2.38; CI = 0.79-2.56). The results of research conducted by Seni Rahayu et al (2019) show that mothers with primiparous and multiparous parity (having less than 4 children) have a lower risk of having stunted toddlers than mothers with

grandemultiparous parity (having more than 4 children) (with OR= 0.4). Parity is an indirect factor in the occurrence of stunting, because parity is closely related to parenting patterns and meeting children's nutritional needs, especially if supported by poor economic conditions. Children born to mothers with high parity have a greater chance of experiencing poor parenting and inadequate nutritional needs during the growth period. Children who have a large number of siblings can cause growth delays due to competition for limited nutritional sources available at home. This is also in line with research conducted by Palino et al (2017) which shows that in the Puuwatu Kendari Health Center working area, toddlers who have mothers with high parity have a 3.25 times greater risk of experiencing stunting compared to toddlers who have mothers with low parity.

CONCLUSION

There was no relationship between the nutritional status of pregnant women, with a V Palue of 0.299, but for the parity variable there was a significant relationship, a V Palue of 0.001.

REFERENCES

- Alsihah (2014). Ekonomi Keluarga dengan Kejadian Stunting anak Usia 24-59 bulan di Wilayah Kerja Puskesmas Barombong Kota Makassar tahun 2014.: *Public Health Science Journal*. 7(1): 63-75
- Annur CM. (2022). Prevalensi Balita Stunting Provinsi Banten Menurut Kabupaten/ Kota. databoks.katadata.co.id/datapublish/2023/02/02/kabupaten-pandeglang-miliki-prevalensi-balita-stunting-tertinggi-di-banten-pada-2022.
- Astutik, Rahfiludin, M. Z., & Aruben, R. (2018). Faktor Risiko Kejadian Stunting pada Anak Balita Usia 24-59 Bulan (Studi Kasus di Wilayah Puskesmas Gabus II Kabupaten Pati Tahun 2017). *Jurnal Kesehatan Masyarakat*, 6(1), 409-418.
- BKKBN, (2021). Buku Kebijakan Dan Strategi Percepatan Penurunan Stunting Di Indonesia.
- Candra, A. 2020. Epidemiologi Stunting. Semarang: Universitas Diponegoro Hubungan Faktor Sosial
- Cholifatun, N., Lailatul, M., (2015). Hubungan Tingkat Pendidikan, Tingkat Pengetahuan dan Pola Asuh Ibu dengan Stunting Pada Pada Balita Keluarga Miskin. *Media Gizi Indonesia*, Vol. 10, No. 1 Januari–Juni: hlm. 84–90
- Enjelia, S.A., Nova, H., Maureen, I., Punuh (2021). Hubungan Antara Tinggi Badan Orangtua dengan Kejadian Stunting pada anak usia 24-59 bulan di kecamatan Tombatu Kabupaten Minahasa Tenggara. *Jurnal Kesehatan Masyarakat, Volume 7 Nomor 4*.
- Febrina, Y., Santoso, S., & Kurniati, A. (2016). No Title. Poltekkes Kemenkes Yogyakarta
- Futihatul, Baidho, (2020). Hubungan Usia, Paritas dan Jarak Kelahiran dengan kejadian Stunting pada balita usia 0-59 di Desa Argodadi
- Kemenkes, R. I. (2018). Situasi Balita Pendek (Stunting) di Indonesia. . Jakarta: Pusat Data dan Informasi, Kementerian, Kesehatan RI.
- Kementerian Kesehatan RI. (2020). Situasi Stunting di Indonesia. Jakarta: Pusat Data dan Informasi Kementerian Kesehatan RI
- Kementrian Kesehatan Republik Indonesia. (2021). Buku Saku Hasil Studi Status Gizi Indonesia (SSGI). <https://www.litbang.kemkes.go.id/buku-saku-hasil-studi-status-gizi-indonesia-ssgi-tahun-2021/>
- Lelemboto, V. S., Malonda, N. S. H., & Punuh, M. I. (2018). Hubungan Antara Tinggi Badan Orangtua Dengan Kejadian Stunting Pada Anak Usia 24-59 Bulan Di Kecamatan Ratahan Timur. *Kesmas*, 7(4).
- Manggala, A.K., Kenwa, K.W., Kenwa, M.M., Sakti, A.A., and Sawitri, A.A. (2018). Risk factors of stunting in children aged 24-59 months. *Paediatrica Indonesiana*. 58(5): 205-212.

- Manuaba. (2013). Ilmu Kebidanan Penyakit Kandungan. dan KB Edisi 2. Jakarta: EGC
- Masturoh, I. dan Anggita, N. (2018). Metodologi Penelitian Kesehatan. Jakarta: Kementerian Kesehatan RI
- Mugianti, dkk, (2018). Faktor penyebab Stunting usia 25-60 bulan di Kecamatan Sukorejo Kota Blitar, *Jurnal Ners dan Kebidanan vol 5 no.3 Desember 2018, hal 268-278.*
- Murti, B. (2003). Prinsip dan Metode Riset Epidemiologi. Yogyakarta: Gadjah Mada University Press
- Murti, F.C., Suryati, dan Oktavianto, E. (2020). Hubungan Berat Badan Lahir Rendah (BBLR) dengan Kejadian Stunting pada Balita usia 2-5 tahun di Desa Umbulrejo, Ponjong, Gunung Kidul. *Jurnal Ilmiah Kesehatan Keperawatan, 16(2): 52-60.*
- Nappu, S., Akri, Y.J., dan Suhartik. (2021). Hubungan Paritas dan Usia Ibu dengan Kejadian BBLR di RS Ben Mari Malang. *Biomed Science. 7(2): 32-42.*
- Nationalnews (2023). <https://www.nasionalnews.id/advertorial/launching-aksi-goceng-arief-jangan-ada-lagi-anak-stunting-di-kota-tangerang/>
- Ni'mah K & Siti Rahayu Nadhiroh. (2016). Faktor yang berhubungan dengan kejadian stunting pada balita. *Media Gizi Indonesia, Vol. 10, No. 1 Januari–Juni 2015: hlm. 13-19.*
- Nisa, L. S., (2018). Kebijakan Penanggulangan Stunting di Indonesia. Stunting Prevention Policies in Indonesia. Badan Penelitian dan Pengembangan Daerah Provinsi Kalimantan Selatan.
- Pahmi, K., Sidrotullah, M., & Ruhardi, A. (2021). Epistemologi Sains Modern Dalam Al-Qur'an, 9. Nusa Tenggara Barat: Natshir Al-Kutub Indonesia. Hal 13-14.
- Palino, I.L., Majid, R., dan Ainurafiq. (2017). Determinan Kejadian Stunting pada Balita Usia 12-59 Bulan di Wilayah Kerja Puskesmas Puuwatu Kota Kendari Tahun 2016. *Jurnal Ilmiah Mahasiswa Kesehatan Masyarakat. 2(6): 1-12.*
- Qoyyimah, A.U., Wintoro, P.D., Hartati, L., Hubungan Riwayat Ibu Hamil Kekurangan Energi Kronis Dengan Kejadian Stunting Pada Balita Usia 3-5 Tahun Di Puskesmas Jatinom Klaten . *Prosiding Seminar Nasional UNIMUS vol.4 2021 1783-1789.*
- Rahayu, A., Yulidasari, F., Putri, A.O., dan Rahman, F. (2015). Riwayat Berat Badan Lahir dengan Kejadian Stunting pada Anak Usia Bawah Dua Tahun. *Jurnal Kesehatan Masyarakat Nasional. 10(2): 67-73.*
- Ramadhan, M.H., Salawati, L., dan Yusuf, S. (2020). Hubungan Tinggi Badan Ibu, Sosial Ekonomi dan Asupan Sumber Zinc Dengan Kejadian Stunting Pada Anak Usia 3-5 Tahun di Puskesmas Kopelma Darussalam. *Jurnal Averrous. 6(1): 55- 65.*
- Ratu, N. C., Punduh, M. I., & Malonda, N. S. (2018). Hubungan tinggi badan orangtua dengan kejadian stunting pada anak usia 24-59 bulan di kecamatan Ratahan kabupaten Minahasa Tenggara. *KESMAS, 7(4).*
- Rifdiani, I. (2016). Pengaruh Paritas, BBL, Jarak Kehamilan, dan Riwayat Perdarahan terjadap Kejadian Perdarahan Postpartum. *Jurnal Berkala Epidemiologi. 4(3): 384-395.*
- Rosselo, J., Kandarina, I., & Kumorowulan, S. (2019). Faktor Risiko Stunting Di Daerah Endemik Gaki Kabupaten Timor Tengah Utara. *Media Gizi Mikro Indonesia, 10(2), 125-136.*
- Sani, M., Solehati, T., & Hendarwati, S. (2020). Hubungan usia ibu saat hamil dengan stunted pada balita 24-59 bulan. *Holistik Jurnal Kesehatan, 13(4), 284–291.* <https://doi.org/10.33024/hjk.v13i4.2016>
- Satriawan. E., (2018). Strategi Nasional Percepatan Pencegahan Stunting 2018-2024. Koordinator Pokja Kebijakan Tim Nasional Percepatan Penanggulangan Kemiskinan (TNP2K) Sekretariat Wakil Presiden Republik Indonesia.

https://www.tnp2k.go.id/filemanager/files/Rakornis%202018/Sesi%201_01_RakorStuntingTNP2K_Stranas_22Nov2018.pdf

- Setiawan, D.A., Suherman, Yusuf, K., Wahyuni, F., dan Vitariani, A. (2021). Faktor yang Berhubungan dengan Kejadian Stunting pada Balita di Puskesmas Ilwaki Kabupaten Maluku Barat Daya Tahun 2021. *PharmaCine: Journal of Pharmacy, Medical and Health Science*. 2(2): 1-14.
- Suhartin, P. (2020). Faktor-Faktor yang Berhubungan dengan Kejadian Stunting Di Kabupaten Konawe Selatan. *Jurnal Ilmiah Kebidanan (Scientific Journal of Midwifery)*, 6 (2), 95-104.
- Sulistyoningsih, H. (2011). Gizi untuk Kesehatan Ibu dan Anak. Yogyakarta: Graha Ilmu.
- Suryani Mokodompit, E. P., Kapantow, N. H., & Mayulu, N. (2019). Hubungan Antara Tinggi Badan Orang Tua Dengan Kejadian Stunting Pada Anak Usia 24-59 Bulan Di Wilayah Kerja Puskesmas Pusomaen Kabupaten Minahasa Tenggara. *KESMAS*, 7(5).
- Susilowati, K. (2016). Gizi Dalam Daur Kehidupan. Bandung: PT Refika Aditama.
- Sutarto, Azqinar, T.A., Himayani, R., dan Wardoyo. (2020). Hubungan Tingkat Pendidikan Ibu dan Pendapatan Keluarga dengan Kejadian Stunting pada Balita di Wilayah Kerja Puskesmas Way Urang Kabupaten Lampung Selatan. *Jurnal Dunia Kesmas*. 9(2): 256-263.
- Syahputra, T. A., Syahrizal and Farizca, A. (2022). Hubungan Antara Kesehatan Mental Ibu dengan Pola Asuh Terhadap Anak', *Jurnal Kedokteran Naggroe Medika*, 5(1), pp. 11–17. doi: 10.24036/perspektif.v4i4.466.
- Thamaria, N. (2017). Penilaian Status Gizi. Jakarta: Kementerian Kesehatan RI
- Tim Nasional Percepatan Penanggulangan Kemiskinan. (2017). 100 Kabupaten/Kota Prioritas untuk Intervensi Anak Kerdil (Stunting). Jakarta: Sekretariat Wakil Presiden Republik Indonesia
- Tri Siswati. (2018). Stunting (pertama). Husada Mandiri Poltekkes Kemenkes Yogyakarta.
- Triwarsini, K., Hadi, H., Siti, N. D, (2016). Riwayat KEK dan anemia pada ibu hamil tidak berhubungan dengan kejadian stunting pada anak usia 6-23 bulan di Kecamatan Sedayu, Bantul, Yogyakarta
- UNICEF, (2017). Malnutrition in Children. [Online]; 2017 (diunduh 23 Maret 2019). Tersedia dari: <https://data.unicef.org/topic/nutrition/malnutrition>.
- Warastuti, Y., & Nengsih, D. (2020). Faktor Risiko Kejadian Stunting Pada Bayi Dan Balita Di Desa Ciambar Kecamatan Ciambar Kabupaten Sukabumi. *Jurnal Kesehatan Dan Kebidanan. Journal Of Health And Midwifery.*, 9 (1), 1-11